

Forage/Haylage Mixing Chart



Directions: Add some water to the applicator tank, then add the required amount of inoculant and continue filling the tank.

- **One Tablespoon of inoculant is 10 grams.**
- **0.7 grams of inoculant treats 1 ton of forage or haylage.**
- When in doubt, always add more inoculant. Extra coverage will not damage the crop.
- Please, see charts below for wagon size.
- These directions do not apply to Pro-Siler or Dohrmann Applicators. See other PDF for Pro-Siler/Dohrmann mixing.

| 6 Ton Wagon Loads | |
|-------------------|--------------------------|
| Number of Loads | Tablespoons of Inoculant |
| 5 | 2¼ |
| 10 | 4½ |
| 15 | 6½ |
| 20 | 8½ |
| 25 | 10½ |

| 8 Ton Wagon Loads | |
|-------------------|--------------------------|
| Number of Loads | Tablespoons of Inoculant |
| 5 | 3 |
| 10 | 5¾ |
| 15 | 8½ |
| 20 | 11¼ |
| 25 | 14 |

| 10 Ton Wagon Loads | |
|--------------------|--------------------------|
| Number of Loads | Tablespoons of Inoculant |
| 5 | 3½ |
| 10 | 7 |
| 15 | 10½ |
| 20 | 14 |
| 25 | 17½ |

| 12 Ton Wagon Loads | |
|--------------------|--------------------------|
| Number of Loads | Tablespoons of Inoculant |
| 5 | 4½ |
| 10 | 8½ |
| 15 | 12¾ |
| 20 | 17 |
| 25 | 21 |

More Info

- Diluting inoculant in a larger volume of water ensures maximum coverage.
- Do not use chlorinated water.
- Do not use hot water.
- For best results use in a sprayer that has a recirculating pump.
- Use only FLOOD jets.
- Use NO SCREENS.
- Use any leftover mixture within 48 hours after mixing.
- Run clear water through applicator after use.
- Store package in a cool dry area.

Any Questions?

Call: 920-251-5916

RK@strongmicrobials.com

StrongMicrobials.com

Made in Milwaukee, WI